

13.Critical Care

These forms are designed to be used by both hospital personnel and external surveyors. The following information must be provided after each survey, before submitting the completed survey forms.

information must be provided after each survey, before submitt	ing the completed survey forms.
1.NAME OF HOSPITAL/CLINIC/FACILITY:	
2. BASELINE/INTERNAL SURVEY INFORMATION:	
Title and name of person who completed this document:	
Post and position held:	
Date of survey:	
3. EXTERNAL SURVEY INFORMATION:	
Name of external surveyor:	
Date of external survey:	
GUIDE TO COMPLETION	ON OF FORM
N.B. Hospital staff are please to use BLACK ink at all time	nes. The external surveyors are requested to
use RED ink at all times.	
Please circle the rated compliance with the criterion, e.g. (Partially compliant), C (Compliant).	NA (Not applicable), NC (Non-compliant), PC
The default category affected is designated on the form to each criterion as follows: 1. patient and staff safety 2. legality 3. patient care 4. efficiency 5. structure 6. basic management 7. basic process 8. evaluation	for
The seriousness of the default is designated on the form for each criterion as follows: 1. mild 2. moderate 3. serious	
4. very serious	
	Documents Checked Surveyor: Surveyor:

© Assessment Instrument Document

Date generated: 24/10/2014 Page 1 of 40

© Assessment Instrument Document

Republic of Botswana

Date generated: 24/10/2014 Page 2 of 40



13.1 Coordination of Patient Care

13.1.1 Standard

Republic of Botswana

During all phases of care, there are qualified individuals responsible for the patient's care.

Standard Intent: The individuals who bear overall responsibility for the patient's care or for a particular phase of care are identified in the patient's record or in a manner that is made known to the personnel.

	Criterion	Comments
		Recommendations
Criterion 13.1.1.1	The individuals responsible	
Critical:	for the patient's care are designated.	
Catg: Basic Management + Efficiency	designated.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.1.1.2	The individuals responsible	
Critical:	for patient care are qualified.	
Catg: Basic Management + Legality		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.1.1.3	The individuals responsible	
Critical:	for patient care are identified and made known to the	
Catg: Basic Management + Efficiency	patient and other personnel.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 3 of 40



13.1.2 Standard

Republic of Botswana

The delivery of services is integrated and coordinated amongst care providers.

Standard Intent: The coordination of patient care depends on the exchange of information between the members of the multidisciplinary/interdisciplinary team. This can be through verbal, written or electronic means as determined by organisational policies. The policies should indicate the appropriate means of communication. Clinical leaders should use techniques to better integrate and coordinate care for their patients (for example, team-delivered care, multi-departmental patient care rounds, combined care planning forums, integrated patient records, case managers). The process for working together will be simple and informal when the patient's needs are not complex.

The patient, family and others are included in the decision process when appropriate.

The patient's record contains a history of all care provided by the multidisciplinary/interdisciplinary team and is made available to all relevant caregivers who are authorised to have access to its content.

	0 % 1	0
	Criterion	Comments
		Recommendations
Criterion 13.1.2.1	The patient's clinical records	
Critical:	are completed according to guidelines determined by the	
Catg: Basic Process + Patient Care	organisation.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.1.2.2	The patients' records are up	
Critical:	to date to ensure the transfer of the latest information between care providers.	
Catg: Basic Process + Patient Care		
Compliance	·	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.1.2.3	Information exchanged	
Critical:	includes a summary of the care provided.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

Page 4 of 40

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.1.2.4 Critical: Catg: Basic Process + Patient Care Compliance	Information exchanged includes the patient's progress.	
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.1.2.5	The author can be identified	
Critical:	for each patient record entry.	
Catg: Basic Process + Legality		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.1.2.6	The date of each patient	
Critical:	record entry can be identified.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.1.2.7	The time of each patient	
Critical:	record entry can be identified.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

Date generated: 24/10/2014 Page 5 of 40

13. Critical Care

13.2 Facilities and Equipment

13.2.1 Standard

Adequate resources are available for the provision of safe care to patients in the ward.

Standard Intent: In order to provide safe patient care, each unit requires adequate resources. The physical facilities required include adequate office accommodation for staff; sluice rooms which are hygienically clean at all times; treatment and dressing rooms; and adequate storage space for clean linen. Cleaning equipment is safely stored in a room or cupboard used for this purpose only. There are adequate toilet and bathing facilities for the number of patients in the ward, as determined by national legislation.

Where there is a neonatal ICU, there is a temperature-controlled nursery and it has:

- suitable bassinettes
- photo-therapy lights
- a panel for viewing babies
- a designated area for preparing infant feeds
- a refrigerator for milk feeds only
- facilities allocated for washing utensils used when preparing infant feeds.

There is adequate lighting and ventilation.

Nurse call systems are available at bedsides and in bathrooms and toilets and are connected to the emergency power supply.

Where there is no piped oxygen and vacuum supply, there are mobile oxygen cylinders and vacuum pumps. All necessary fittings for oxygen and suction are in place and working satisfactorily. Each ward is provided with a socket outlet that is connected to the emergency power supply.

A resuscitation trolley is available at the point of need within one minute. In addition, there is access to a defibrillator or automated external defibrillator (AED) within three minutes of any patient collapsing. Resuscitation equipment includes at least:

- a defibrillator with adult paddles/pads (and infant paddles/pads where applicable)
- an ECG monitor
- a CPR board (if required)
- suction apparatus (electrical or alternative) plus a range of soft and hard suction
- a bag-mask manual ventilator
- a range of endotracheal tubes and two laryngoscopes with a range of straight and curved blades, spare batteries, spare globes where applicable
- an introducer/stylet for endotracheal intubation
- a syringe to inflate the ETT cuff
- oropharyngeal tubes
- equipment to perform an emergency cricothyroidotomy (needle and surgical)
- appropriate facilities for intravenous therapy and drug administration (including
- drugs for cardiac arrest, coma, seizures and states of shock (including paediatric doses where applicable)
- plasma expanders
- pulse oximeter.

Criterion	Comments	
	Recommendations	

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.2.1.1 Critical: Catg: Basic Management + Physical Struct Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Patient and staff accommodation and equipment is adequate to meet patient care needs.	
Criterion 13.2.1.2 Critical: Catg: Basic Management + Physical Struct Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Oxygen and vacuum supplies meet the needs of patients for care.	
Criterion 13.2.1.3 Critical: Catg: Basic Process + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	There is evidence that equipment is maintained in accordance with the policies of the organisation. 7	
Criterion 13.2.1.4 Critical: Catg: Basic Management + Physical Struct Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Resuscitation equipment is available in accordance with the policies of the organisation.	
Criterion 13.2.1.5 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Where there are no piped oxygen installations, there is a documented procedure for ensuring that cylinder pressures (i.e. contents) are monitored according to organisational policy while patients are receiving oxygen.	

© Assessment Instrument Document

Date generated: 24/10/2014

Page 7 of 40

Republic of Botswana

13.Critical Care

Criterion 13.2.1.6	Each patient has access to a	
Critical:	nurse call system at all times.	
Catg: Basic Management + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.2.1.7	Electricity and water is	
Critical:	available in accordance with the policies of the	
Catg: Basic Management + Physical Struct	organisation.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.2.1.8	Where there is a neonatal	
Critical:	ICU, there is a dedicated area for preparing infant feeds.	
Catg: Basic Management + Physical Struct	for proparing illiant roods.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

13.2.2 Standard

Specific resources are available for the provision of safe care to patients in the critical care unit.

Standard Intent: Professional guidelines for critical care services recommend the personnel and resources required to manage the service safely. The personnel in the ward are in possession of these guidelines and ensure that the recommendations are implemented. These guidelines provide norms for staffing a critical care unit and also for the services and facilities required.

	Criterion	Comments
		Recommendations
Criterion 13.2.2.1	Current guidelines for the	
Critical:	provision of critical care services and facilities are	
Catg: Basic Management + Efficiency	followed.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 8 of 40

Republic of Botswana

13.Critical Care

Criterion 13.2.2.2 Critical: Catg: Basic Management + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Staffing of the service complies with accepted staffing norms for critical care services.	
Criterion 13.2.2.3 Critical: Catg: Basic Management + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Available medical equipment complies with accepted norms for critical care services.	
Criterion 13.2.2.4 Critical: Catg: Basic Process + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Where resuscitation, intensive care, life support or critical monitoring equipment is used that does not have built-in battery backup units, there is an uninterruptible power supply (UPS) that complies with relevant requirements and is regularly serviced and tested.	

Date generated: 24/10/2014 Page 9 of 40



13.3 Clinical Practice Guidelines

13.3.1 Standard

Republic of Botswana

Clinical practice guidelines are used to guide patient care and reduce unwanted variation.

Standard Intent: Clinical practice guidelines provide a means for improving quality and they assist practitioners and patients in making clinical decisions. Guidelines are found in the literature under many names, including practice parameters, practice guidelines, patient care protocols, standards of practice and/or care pathways. Regardless of the source, the scientific basis of guidelines should be reviewed and approved by organisational leaders and clinical practitioners before implementation. Consideration should be given to providing guidelines for high risk, high volume and high cost conditions as these will form the basis for structured clinical audits.

This ensures that they meet the criteria established by the leaders and are adapted to the community, patient needs and organisational resources. Once implemented, guidelines are reviewed on a regular basis to ensure their continued relevance.

	Criterion	Comments
		Recommendations
Criterion 13.3.1.1 Critical: Catg: Basic Process + Pat & Staff Safety Compliance	Clinical practice guidelines relevant to the patients and services of the organisation are available to guide patient care processes.	
NA NC PC C Default Severity for NC or PC = 4 Very Serious		
Criterion 13.3.1.2	The implementation of	
Critical:	guidelines is monitored as part of a structured clinical	
Catg: Evaluation + Patient Care	audit.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.3.1.3	Guidelines are reviewed and adapted on a regular basis.	
Catg: Evaluation + Efficiency Compliance		
NA NC PC C Default Severity for NC or PC = 3 Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 10 of 40



13.4 Assessment of Patients

13.4.1 Standard

Republic of Botswana

All patients cared for by the organisation have their health needs identified through an established assessment process.

Standard Intent: When a patient enters a ward, the specific information required and the procedures for obtaining and documenting it depend on the patient's needs and on the setting in which care is being provided.

The organisation defines in writing the scope and content of assessments to be performed by each clinical discipline within its scope of practice and applicable laws and regulations. These findings are used throughout the care process to evaluate patient progress and provide information regarding the need for re-assessment. It is essential that assessments are well documented and can be easily retrieved from the patient's record.

The health organisation determines the time frame for completing assessments. This may vary in the different settings within the organisation. When an assessment is partially or entirely completed outside the organisation, the findings are verified on admission to the organisation.

	Criterion	Comments
		Recommendations
Criterion 13.4.1.1	The organisation implements policies and procedures for	
Catg: Basic Management + Patient Care	assessing patients on admission and during on-	
Compliance	going care.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.4.1.2	Only those individuals	
Critical:	permitted by applicable laws and regulations or by	
Catg: Basic Process + Legality	registration perform the assessments.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.4.1.3	The scope and content of	
Critical:	assessment by each discipline is defined.	
Catg: Basic Management + Patient Care	uiscipiirie is defined.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 11 of 40



13. Critical Care

Criterion 1	3.4.1.4			Policies and procedures	
Critical:				ensure that assessments are performed within appropriate	
Catg: Basic	c Manag	ement -	+ Patient	time frames and that they are adequately documented in	
	Compl	iance		the patient records.	
NA	NC	PC	С		
Default Sev Serious	verity for	NC or I	PC = 3		

13.4.2 Standard

Each patient has an initial assessment that complies with current policies, procedures and guidelines.

Standard Intent: The initial assessment of a patient is critical for the identification of the needs of the patient and initiation of the care process. A patient's social, cultural and family status are important factors that can influence their response to illness and care. Families can be of considerable help in these areas of assessment and in understanding the patient's wishes and preferences. Economic factors are assessed as part of the social assessment, particularly when the patient and his/her family will be responsible for the cost of all or a portion of the care.

A functional and nutritional assessment allows for the patient to be referred for specialist care if necessary.

Certain patients may require a modified assessment, e.g. very young patients, the frail or elderly, those terminally ill or in pain, patients suspected of drug and/or alcohol dependence and victims of abuse and neglect. The assessment process is modified in accordance with national guidelines and protocols. The outcome from the patient's initial assessment results in an understanding of the patient's medical and nursing needs so that care and treatment can begin.

Planning for discharge is initiated during the initial assessment process.

When the medical assessment was conducted outside the organisation, a legible copy of the findings is placed in the patient's record. Any significant changes in the patient's condition since this assessment are recorded.

				Criterion	Comments
					Recommendations
Criterion 13.4.2.1				Each patient admitted has an	
Critical:				initial assessment that meets organisational policy.	
Catg: Basic Process + Patient Care			ient Care		
Compliance					
NA	NC	PC	С		
Default Severity for NC or PC = 4 Very Serious			PC = 4		

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.4.2.2	The initial assessment includes health history.	
Critical:	 	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.4.2.3	The initial assessment	
Critical:	includes physical examination.	
Catg: Basic Process + Patient Care	examination.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.4.2.4	The initial assessment	
Critical:	includes functional examination, where applicable.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.4.2.5	The initial assessment	
Critical:	includes social and economic	
Catg: Basic Process + Patient Care	assessment, where applicable.	
Compliance	` `	
NA NC PC C		
Default Severity for NC or PC = 2 Moderate		
Criterion 13.4.2.6	The initial assessment	
Critical:	includes psychological assessment, where applicable.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 2 Moderate		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 13 of 40

Republic of Botswana

13.Critical Care

Criterion 13.4.2.7 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	The initial assessment includes cultural assessment, where applicable.	
Criterion 13.4.2.8 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	The initial assessment results in an initial diagnosis.	
Criterion 13.4.2.9 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	The initial assessment results in the identification of the patient's medical, nursing or other health needs.	

13.4.3 Standard

Health professionals responsible for patient care collaborate to analyse and integrate assessment information.

Standard Intent: A patient benefits most when the personnel responsible for the patient work together to analyse the assessment findings and to combine this information into a comprehensive picture of his or her condition. From this collaboration, the patient's needs are identified, the order of their importance is established and care decisions are made.

	Criterion	Comments
		Recommendations
Criterion 13.4.3.1	Assessment findings are	
Critical:	documented in the patient's record and are readily	
Catg: Basic Process + Patient Care	available to those responsible	
Compliance	for the patient's care.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 14 of 40

Republic of Botswana

13.Critical Care

Criterion 13.4.3.2 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Patient assessment data and information are analysed and integrated by those responsible for the patient's care.	
Criterion 13.4.3.3 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Patient needs are prioritised on the basis of assessment results.	
Criterion 13.4.3.4 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	The patient and/or the family participate in the decisions regarding the priority needs to be met.	

Date generated: 24/10/2014 Page 15 of 40

13.5 Patient Care

Republic of Botswana

13.5.1 Standard

The care provided to each patient is planned and written in the patient's record.

Standard Intent: A single integrated plan is preferable to a separate care plan recorded by each health professional.

Collaborative care and treatment team meetings or similar patient discussions are recorded.

Individuals qualified to do so order diagnostic and other procedures. These orders must be easily accessible if they are to be acted on in a timely manner. Locating orders on a common sheet or in a uniform location in patient records facilitates the correct understanding and carrying out of orders.

The organisation decides:

- which orders must be written rather than verbal
- who is permitted to write orders
- where orders are to be located in the patient record.

The method used must respect the confidentiality of patient care information.

	Criterion	Comments
		Recommendations
Criterion 13.5.1.1	The planned care is provided and noted in the patient's record.	
Critical:		
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.1.2	All procedures and diagnostic	
Critical:	tests ordered and performed are written into the patient's record.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.1.3	The results of procedures and	
Critical:	diagnostic tests performed are available in the patient's record.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.5.1.4	Re-assessments are documented in the patient's record.	
Catg: Basic Process + Patient Care Compliance NA NC PC C	Todord.	
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.1.5	The patient's plan of care is modified when the patient's needs change.	
Catg: Basic Process + Patient Care Compliance	noods ondings.	
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

Date generated: 24/10/2014 Page 17 of 40



13.5.2 Standard

Republic of Botswana

Policies and procedures guide the care of high-risk patients and the provision of high-risk services.

Standard Intent: Some patients are considered "high-risk" because of their age, condition or the critical nature of their needs. Children and the elderly are commonly in this group as they may not be able to speak for themselves, understand the care process or participate in decisions regarding their care. Similarly, the frightened, confused or comatose patient is unable to understand the care process when care needs to be provided efficiently and rapidly.

Policies and procedures are important. They help the personnel understand these patients and services and respond in a thorough, competent and uniform manner. The clinical and managerial leaders take responsibility for identifying the patients and services considered high-risk, using a collaborative process to develop policies and procedures and train staff in their implementation.

The special facilities and safety measures required by children need to be specified.

It is particularly important that the policies or procedures indicate:

- how planning will occur
- the documentation required for the care team to work effectively
- special consent considerations
- monitoring requirements
- special qualifications or skills of the personnel involved in the care process and
- the resuscitation equipment available and how to use it, including equipment for children.

Clinical guidelines should be incorporated in the process because there are several criteria requiring guidelines to be used. Monitoring provides the information needed to ensure that the policies and procedures are adequately implemented and followed for all relevant patients and services.

Policies and procedures should focus on high-risk patients and procedures, e.g.

- a) the care of emergency patients
- b) the handling, use and administration of blood and blood products
- c) the management of contaminated blood supplies (expired, opened or damaged container)
 d) the care of patients on life support or those who are comatose
- e) the care of patients with communicable diseases
- f) the care of immuno-suppressed patients
- g) the care of patients on dialysis
- h) the use of restraint and the care of patients in restraint
- i) the care of frail, dependent, elderly patients
- j) the care of young, dependent children
- k) the security of newborn babies, where applicable
- I) the care of adolescents.

	Criterion	Comments
		Recommendations
Criterion 13.5.2.1	Policies and procedures for	
Critical:	identified high-risk patients and procedures, which	
Catg: Basic Process + Patient Care	include at least items a) to l)	
Compliance	in the intent statement above, are implemented.	
NA NC PC C	are implemented.	
Default Severity for NC or PC = 4 Very Serious		



Republic of Botswana 13. Critical Care

				The personnel are trained	
				and use the policies and procedures to guide care.	
Catg: Basic Process + Patient Care			tient Care		
Compliance					
NA	NC	PC	С		
Default Severity for NC or PC = 3 Serious			PC = 3		

13.5.3 Standard

Risks, benefits, potential complications, and care options are discussed with the patient and his or her family or with those who make decisions for the patient.

Standard Intent: This section deal with the process of obtaining informed consent from patients and does NOT refer to providing health education in general, which is dealt with in Standard 13.8.

Patients and their families or decision-makers receive adequate information to participate in care decisions. Patients and families are informed as to what tests, procedures and treatments require consent and how they can give consent. For example, consent may be given verbally, by signing a consent form or through some other mechanism. Patients and families understand who may give consent in addition to the patient.

Designated personnel are trained to inform patients and to obtain and document patient consent, e.g. a doctor for a surgical procedure or a nurse for HIV testing. These staff members clearly explain any proposed treatments or procedures to the patient and, when appropriate, the family. Informed consent includes:

- an explanation of the risks and benefits of the planned procedure
- identification of potential complications
- consideration of the surgical and non-surgical options available to treat the patient.

In addition, when blood or blood products may be needed information on the risks and alternatives is discussed.

The organisation lists all those procedures that require written informed consent. Leaders document the processes for obtaining informed consent.

The consent process always concludes with the patient signing the consent form, or the documentation of the patient's verbal consent in the patient's record by the individual who provided the information for consent. Documentation includes the statement that the patient acknowledges full understanding of the information. The patient's surgeon or other qualified individual provides the necessary information and the name of this person appears on the consent form.

	Criterion	Comments	
		Recommendations	
Criterion 13.5.3.1	There is a documented		
Critical:	process for obtaining informed consent.		
Catg: Basic Management + Patient Care			
Compliance			
NA NC PC C			
Default Severity for NC or PC = 4 Very Serious			

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.5.3.2 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C	Patients are informed about their condition and the proposed treatment.	
Default Severity for NC or PC = 3 Serious		
Criterion 13.5.3.3 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Patients know the identity of the medical practitioner or other professional practitioner responsible for their care.	
Criterion 13.5.3.4 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	The information provided is recorded with the record of the patient having provided written or verbal consent.	

Date generated: 24/10/2014 Page 20 of 40



13.5.4 Standard

Republic of Botswana

Pre- and post-operative assessments are documented.

Standard Intent: The pre-operative anaesthetic assessment determines whether the patient is a good candidate for the planned surgery and may significantly influence the pre- and intra-operative management. The clinical assessment and results of investigations must be available to the doctor performing the assessment. In an emergency, the initial medical assessment may be limited to the patient's apparent needs and condition. Appropriate re-assessments are essential to modify and guide effective treatment.

A patient's post-surgical care is related to the findings and the surgical procedure. The surgical report is available within a time frame needed to provide post-surgical care to the patient. Post-operative monitoring is appropriate to the patient's condition and the procedure performed.

Results of monitoring influence intra- and post-operative decisions such as return to surgery, transfer to another level of care and the need for further investigations or discharge.

	Criterion	Comments
		Recommendations
Criterion 13.5.4.1	The patient's initial medical	
Critical:	assessment is documented before anaesthesia.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.4.2	The patient's pre-operative	
Critical:	diagnosis is recorded before anaesthesia.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.4.3	A post-operative diagnosis is	
Critical:	documented.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 21 of 40



13.Critical Care

Criterion 13.5.4.4	The name of the surgeon and	
Critical:	the names of other personnel as required by law are	
Catg: Basic Process + Legality	documented.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.4.5	The patient's physiological	
Critical:	status is monitored during the immediate post-surgery	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

13.5.5 Standard

The organisation implements processes to support the patient in pain management.

Standard Intent: While pain may be a part of the patient experience, unrelieved pain has adverse physical and psychological effects. The patient's right to appropriate assessment and management of pain is respected and supported.

The organisation has processes to:

- identify patients with pain during initial assessment and re-assessment
- communicate with and provide education for patients and families about pain management in the context of their personal, cultural and religious beliefs
- educate health service providers in pain assessment and management.

	Criterion	Comments
		Recommendations
Criterion 13.5.5.1	The assessment process	
Critical:	makes provision for patients in pain to be identified.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.5.2	Patients in pain receive care	
Critical:	according to pain management guidelines.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 22 of 40

Republic of Botswana

13.Critical Care

Criterion 13.5.5.3	Patients and families are educated about pain management.	
Catg: Basic Process + Patient Care Compliance NA NC PC C	management.	
Default Severity for NC or PC = 3 Serious		
Criterion 13.5.5.4 Critical: Catg: Basic Management + Efficiency	The organisation has processes to educate health professionals in assessing and managing pain.	
Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious		

13.5.6 Standard

The organisation develops processes to manage end-of-life care.

Standard Intent: Dying patients have unique needs for respectful, compassionate care. Concern for the patient's comfort and dignity guides all aspects of care during the final stages of life. To accomplish this, all personnel are made aware of the unique needs of patients at the end of life. These needs include treatment of primary and secondary symptoms, pain management, responding to the concerns of the patient and their family and involving them in care decisions.

End-of-life care provided by the organisation includes:

- a) providing appropriate treatment for any symptoms according to the wishes of the patient and family
- b) sensitively addressing issues such as autopsy and organ donation
- c) involving the patient and family in all aspects of care
- d) responding to the psychological, emotional, spiritual and cultural concerns of the patient and family.

	Criterion	Comments
		Recommendations
Criterion 13.5.6.1	Policies and procedures	
Critical:	regarding end-of-life care, at least including elements a) to	
Catg: Basic Process + Patient Care	d) in the intent statement, are	
Compliance	implemented.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 23 of 40

Republic of Botswana

13.Critical Care

Criterion 13.5.6.2 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	The patient and the family/guardian are involved in care decisions.	
Serious		
Criterion 13.5.6.3	Pain and primary or secondary symptoms are managed.	
Catg: Basic Process + Patient Care Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.5.6.4	Interventions address patient	
Critical:	and family religious and cultural concerns.	
Catg: Basic Process + Patient Care	Cultural Concerns.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

Date generated: 24/10/2014 Page 24 of 40



13.Critical Care

13.6 Medication

13.6.1 Standard

Medication use in the organisation complies with applicable laws and regulations.

Standard Intent: Medication management is not only the responsibility of the pharmaceutical service but also of managers and clinical care providers. Medical, nursing, pharmacy and administrative personnel participate in a collaborative process to develop and monitor policies and procedures.

Each organisation has a responsibility to identify those individuals with the requisite knowledge and experience, and who are permitted by law, registration or regulations to prescribe or order medications. In emergency situations, the organisation identifies any additional individuals permitted to prescribe or order medications. Requirements for documentation of medications ordered or prescribed and for using verbal medication orders are defined in policy.

	Criterion	Comments
		Recommendations
Criterion 13.6.1.1	Policies and procedures that	
Critical:	guide the safe prescribing, ordering, storage and	
Catg: Basic Management + Legality Compliance	administration of medications are implemented.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.6.1.2	The use of verbal/telephonic	
Critical:	medication orders is documented.	
Catg: Basic Management + Legality	accumented.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.6.1.3	Only those permitted by the	
Critical:	organisation and by relevant laws and regulations	
Catg: Basic Process + Legality	prescribe medication.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		



13.Critical Care

Criterion 13.6.1.4	Medications, including herbal	
Critical:	and ove- the-counter medications, brought into the	
Catg: Basic Process + Patient Care	organisation by the patient or	
Compliance	the family are known to the patient's medical practitioner	
NA NC PC C	and are noted in the patient's	
Default Severity for NC or PC = 3 Serious	record.	

13.6.2 Standard

Medications are safely administered.

Standard Intent: Only personnel who are suitably trained and experienced may administer medication to patients. The responsibility of these persons for medication administration is documented. The safe administration of medications requires a strict and comprehensive protocol.

The patient, medical practitioner, nurse and other care providers work together to monitor patients on medications. The purpose of monitoring is to evaluate the response to medication, adjust the dosage or type of medication when needed and to evaluate the patient for adverse effects.

The organisation follows national requirements for the reporting of adverse effects. Doctors, nurses and pharmacists are expected to report reactions that are suspected to be adverse drug events, irrespective of whether the event is well recognised, potentially serious or clinically "insignificant".

There is a reporting process focused on the prevention of medication errors through understanding the types of errors that occur. Improvements in medication processes and staff training are used to prevent errors in the future. The pharmacy participates in such staff training.

	Criterion	Comments
		Recommendations
Criterion 13.6.2.1	Only those permitted by the	
Critical:	organisation and by relevant laws and regulations	
Catg: Basic Process + Legality	administer medications.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.6.2.2	There is evidence that	
Critical:	patients are identified before medications are	
Catg: Basic Process + Pat & Staff Safety	administered.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 26 of 40

Republic of Botswana

13.Critical Care

Criterion 13.6.2.3 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Medications are checked against the original prescriptions and administered as prescribed.	
Criterion 13.6.2.4 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Health professionals monitor medication effects on patients collaboratively.	
Criterion 13.6.2.5 Critical: Catg: Basic Process + Legality Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Adverse Drug Reactions (ADR) are observed, recorded and reported through a process and within a time frame defined by the organisation.	
Criterion 13.6.2.6 Critical: Catg: Basic Process + Pat & Staff Safety Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Medication errors are reported through a process and within a time frame defined by the organisation.	
Criterion 13.6.2.7 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	The medications prescribed for and administered to each patient are recorded.	

© Assessment Instrument Document

Date generated: 24/10/2014 Page 27 of 40



Republic of Botswana 13. Critical Care

13.6.3 Standard

Medications are stored in a safe and clean environment.

Standard Intent: Patient care units store medications in a clean and safe environment that complies with law, regulation and professional practice standards.

	Criterion	Comments
		Recommendations
Criterion 13.6.3.1	Medication is stored in a	
Critical:	locked storage device or cabinet that is accessible only	
Catg: Basic Process + Efficiency	to authorised personnel.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.6.3.2	Medications identified for	
Critical:	special control (by law or or organisational policy) are	
Catg: Basic Process + Legality	stored in a cabinet of	
Compliance	substantial construction, for	
NA NC PC C	which only authorised personnel have the keys.	
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.6.3.3	Medications identified for	
Critical: D	special control (by law or or organisational policy) are	
Catg: Basic Process + Legality	accurately accounted for.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.6.3.4	Medications are securely and	
Critical: D	legibly labelled with relevant information as required by	
Catg: Basic Process + Pat & Staff Safety	law and organisational policy.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 28 of 40

Republic of Botswana

13.Critical Care

Criterion 13.6.3.5 Critical: Catg: Basic Process + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Medications are stored in a clean environment.	
Criterion 13.6.3.6 Critical: Catg: Basic Management + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Medication is stored in accordance with manufacturer's instructions relating to temperature, light and humidity.	
Criterion 13.6.3.7 Critical: Catg: Basic Management + Physical Struct Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	A lockable refrigerator is available for those medications requiring storage at low temperatures.	
Criterion 13.6.3.8 Critical: O Catg: Basic Process + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	The temperature of the refrigerator is monitored and recorded.	
Criterion 13.6.3.9 Critical: Catg: Basic Process + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Expiry dates are checked (including those of emergency drugs), and drugs are replaced before expiry date.	

© Assessment Instrument Document

Date generated: 24/10/2014 Page 29 of 40



13.7 Food and Nutrition Therapy

13.7.1 Standard

Republic of Botswana

Food and nutrition therapy appropriate for the patient and consistent with his or her clinical care is regularly available.

Standard Intent: A qualified caregiver orders appropriate food or other nutrients. The patient participates in planning and selecting foods and the patient's family may, when appropriate, participate in providing food. They are educated as to which foods are contraindicated, including information about any medications associated with food interactions. When possible, patients are offered a variety of food choices consistent with their nutritional status.

The nutritional status of the patients is monitored.

	Criterion	Comments
		Recommendations
Criterion 13.7.1.1	Food, appropriate to the	
Critical:	patient, is regularly available.	
Catg: Basic Management + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.7.1.2	An order for food based on	
Critical:	the patient's nutritional status and needs is recorded in the	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.7.1.3	When families provide food,	
Critical:	they are educated about the patient's diet limitations.	
Catg: Basic Process + Patient Care	pation o diot infinations.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.7.1.4	Patients assessed as being at	
Critical:	nutrition risk receive nutrition therapy.	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.7.1.5 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	A collaborative process is used to plan, deliver and monitor nutrition therapy.	
Criterion 13.7.1.6 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 3 Serious	Nutrition therapy provided, either oral or intravenous, is written in the patient's record.	
Criterion 13.7.1.7 Critical: Catg: Basic Process + Patient Care Compliance NA NC PC C Default Severity for NC or PC = 4 Very Serious	Response to nutrition therapy is monitored and recorded.	

Date generated: 24/10/2014 Page 31 of 40

13.8 Patient and Family Education

13.8.1 Standard

Republic of Botswana

Education supports patient and family participation in care decisions and processes.

Standard Intent: Learning occurs when attention is paid to the methods used to educate patients and families. The organisation selects appropriate educational methods and people to provide the education.

Staff collaboration helps to ensure that the information patients and families receive is comprehensive, consistent and as effective as possible.

Education is focused on the specific knowledge and skills that the patient and his or her family will need to make care decisions, participate in care and continue care at home. Variables like educational literacy, beliefs and limitations are taken into account.

Each organisation decides on the placement and format for educational assessment, planning and delivery of information in the patient's record.

Education is provided to support care decisions of patients and families. In addition, when a patient or family directly participates in providing care, for example changing dressings, feeding and administration, they need to be educated.

It is sometimes important that patients and families are made aware of any financial implications associated with care choices, such as choosing to remain an inpatient rather than being an outpatient.

Education in areas that carry high risk to patients is routinely provided by the organisation, for instance instruction in the safe and effective use of medications and medical equipment. Community organisations that support health promotion and disease prevention education are identified and, when possible, on-going relationships are established.

	Criterion	Comments
		Recommendations
Criterion 13.8.1.1	Patients and families indicate	
Critical:	that they have been informed about their diagnosis.	
Catg: Basic Process + Patient Care	and at anom anagricoid.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 2 Moderate		
Criterion 13.8.1.2	Patients indicate that they	
Critical:	have been informed about the management of their	
Catg: Basic Process + Patient Care	condition.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 2 Moderate		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 32 of 40

Republic of Botswana

13.Critical Care

Criterion 13.8.1.3	Patients are educated about	
Critical:	their diagnosis, relevant high health risks, e.g. safe use of medication and medical equipment, medicine and food interaction, diet and food interactions, defaulting on	
Catg: Basic Process + Patient Care Compliance NA NC PC C		
Default Severity for NC or PC = 4 Very Serious	medication use, etc.	
Criterion 13.8.1.4	Patients and families indicate	
Critical:	that they have been informed about any financial	
Catg: Basic Process + Patient Care	implications of care decisions.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 2 Moderate		

13.9 Continuity of Care

13.9.1 Standard

The organisation designs and carries out processes to provide continuity of patient care services within the organisation and coordination among health professionals.

Standard Intent: As patients move through a health organisation from admission to discharge or transfer, several departments and services and many different health service providers may be involved in providing care. Without coordination and effective transfer of information and responsibilities, errors of omission and commission may occur, exposing the patient to avoidable risks.

	Criterion	Comments	
		Recommendations	
Criterion 13.9.1.1	Established criteria or policies		
Critical:	and procedures that guide the movement of patients within the organisation are implemented.		
Catg: Basic Management + Patient Care			
Compliance	'		
NA NC PC C			
Default Severity for NC or PC = 3 Serious			
Criterion 13.9.1.2	Individuals responsible for the		
Critical:	patient's care and its coordination are identified for all phases.		
Catg: Basic Management + Patient Care			
Compliance			
NA NC PC C			
Default Severity for NC or PC = 3 Serious			

© Assessment Instrument Document

Date generated: 24/10/2014 Page 33 of 40

Republic of Botswana

13.Critical Care

Criterion 13.9.1.3	Continuity and coordination are evident throughout all phases of patient care.	
Critical:		
Catg: Basic Process + Patient Care Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		
Criterion 13.9.1.4	The record of the patient accompanies the patient when transferred within the	
Critical:		
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 3 Serious		

13.9.2 Standard

There is a process to appropriately transfer patients to another organisation to meet their continuing needs.

Standard Intent: Transfer may be for specialised consultation and treatment, urgent services or for less intensive services such as sub-acute care or long-term rehabilitation.

To ensure continuity of care, adequate information must accompany the patient.

Transfer may be a brief process with the patient alert and talking or it may involve continuous nursing or medical supervision. The process for transferring the patient must consider transportation needs. The qualifications of the individual accompanying the patient must be appropriate.

	Criterion	Comments
		Recommendations
Criterion 13.9.2.1	There is a documented process for transferring patients to other	
Critical:		
Catg: Basic Management + Patient Care	l'	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.9.2.2	The referring organisation	
Critical:	determines that the receiving organisation can meet the patient's continuing care needs and establishes arrangements to ensure	
Catg: Basic Process + Patient Care		
Compliance		
NA NC PC C	continuity.	
Default Severity for NC or PC = 3 Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 34 of 40

Republic of Botswana

13.Critical Care

The process for transferring	
transportation needs.	
The process determines that	
appropriately qualified person	
during transfer.	
When a patient is transferred	
organisation.	
A copy of the transfer	
summary is available in the	
patient record.	
]	
The health organisation	
agreeing to receive the	
patient's record.	
	The process determines that patients are accompanied and monitored by an appropriately qualified person during transfer. When a patient is transferred to another organisation, the receiving organisation is given a written summary of the patient's clinical condition and the interventions provided by the referring organisation. A copy of the transfer summary is available in the patient record. The health organisation agreeing to receive the patient is noted in the

© Assessment Instrument Document

Date generated: 24/10/2014 Page 35 of 40

13.10 Quality Improvement

13.10.1 Standard

Republic of Botswana

A formalised proactive quality improvement approach is maintained in the service.

Standard Intent: This refers to the implementation of organisational quality improvement processes (Service Element 8).

It is the responsibility of management of the organisation to ensure that standards are set throughout the organisation. Within each department or service, it is the responsibility of managers to ensure that standards are set for the particular department. This requires coordination with the organisation's central, management or coordinating quality improvement structures or systems. Departmental managers use available data and information to identify priority areas for quality monitoring and improvement.

Quality monitoring could include:

- a) patient assessment
- b) surgical procedures carried out
- c) the use of antibiotics and other medications and medication errors
- d) the use of anaesthesia
- e) the use of blood and blood products
- f) patient and family expectations and satisfaction.

The following will be evaluated:

- problems identified in this service for which quality improvement activities were initiated
- the processes put in place to resolve the problems
- the identification of indicators to measure improvement
- the tool(s) used to evaluate these indicators
- the monitioring of these indicators and corrective steps taken when goals were not achieved
- graphed and/or tabled results, as appropriate.

	Criterion	Comments
		Recommendations
Criterion 13.10.1.1	There are formalised quality	
Critical:	improvement processes for the service that have been	
Catg: Evaluation + Efficiency	developed and agreed upon	
Compliance	by the personnel of the service.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.10.1.2	Indicators of performance are	
Critical:	identified to evaluate the quality of treatment and patient care.	
Catg: Evaluation + Efficiency		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Republic of Botswana

13.Critical Care

Criterion 13.10.1.3 Critical: Catg: Evaluation + Efficiency Compliance NA NC PC C Default Severity for NC or PC = 4	The quality improvement cycle includes the monitoring and evaluation of the standards set and the remedial action implemented.	
Very Serious	A de como contesta a constitu	
Criterion 13.10.1.4	A documentation audit system is in place.	
Catg: Evaluation + Efficiency Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

13.11 Patient Rights

13.11.1 Standard

The department/service implements processes that support patient and family rights during care.

Standard Intent: This refers to the implementation of organisational policies on patient and family rights (Service Element 5). Compliance will be verified during observation of patient care processes, patient record audits and patient interviews.

	Criterion	Comments
		Recommendations
Criterion 13.11.1.1	There are processes that	
Critical:	support patient and family rights during care.	
Catg: Basic Management + Patient Care		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.11.1.2	Measures are taken to protect	
Critical:	the patient's privacy, person and possessions.	
Catg: Basic Process + Patient Care	and possessions.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 37 of 40



13.Critical Care

Criterion 1	3.11.1.3	3		The personnel respect the	
			rights of patients and families to treatment and to refuse		
Catg: Basic	Catg: Basic Process + Patient Care		tient Care		
Compliance					
NA	NC	PC	С		
Default Severity for NC or PC = 4 Very Serious		PC = 4			

13.12 Prevention and Control of Infection

13.12.1 Standard

The department/service implements infection prevention and control processes.

Standard Intent: This refers to the implementation of organisational processes for infection prevention and control (Service Element 9).

	Criterion	Comments
		Recommendations
Criterion 13.12.1.1	The department identifies the procedures and processes associated with the risk of infection and implements strategies to reduce risk.	
Catg: Basic Process + Pat & Staff Safety		
Compliance	onatographic to rouges man.	
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.12.1.2	Infection control processes	
Critical:	include prevention of the	
Catg: Basic Process + Pat & Staff Safety	spread of respiratory tract infections.	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.12.1.3	Infection control processes	
Critical:	include prevention of the spread of urinary tract infections.	
Catg: Basic Process + Pat & Staff Safety		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 38 of 40



13.Critical Care

Criterion 13.12.1.4 Critical: Catg: Basic Process + Pat & Staff Safety Compliance NA NC PC C Default Severity for NC or PC = 4	Infection control processes include prevention of the spread of infection through intravascular invasive devices.	
Very Serious		
Criterion 13.12.1.5 Critical: Catg: Basic Process + Pat & Staff	Infection control processes include prevention of the spread of infection through surgical wounds.	
Safety	J g	
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

13.13 Risk Management

13.13.1 Standard

The department/service implements risk management processes.

Standard Intent: This refers to the implementation of organisational risk management processes. (Service Element 7).

	Criterion	Comments
		Recommendations
Criterion 13.13.1.1	The department conducts on-	
Critical:	going monitoring of risks through documented	
Catg: Basic Process + Efficiency	assessments as part of	
Compliance	organisational risk management processes.	
NA NC PC C	·	
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.13.1.2	A system for monitoring incidents, near misses, sentinel and adverse events is available and includes the documentation of interventions and responses to recorded incidents	
Critical:		
Catg: Basic Process + Pat & Staff Safety		
Compliance		
NA NC PC C	ito recorded incidents	
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 39 of 40

Republic of Botswana

13.Critical Care

Criterion 13.13.1.3	Security measures are in place and are implemented to ensure the safety of patients, personnel and visitors.	
Critical:		
Catg: Basic Process + Pat & Staff Safety		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.13.1.4	Fire safety measures are	
Critical:	implemented.	
Catg: Basic Process + Pat & Staff Safety		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		
Criterion 13.13.1.5	The organisation's policy on	
Critical:	handling, storing and disposing of health waste is implemented.	
Catg: Basic Process + Pat & Staff Safety		
Compliance		
NA NC PC C		
Default Severity for NC or PC = 4 Very Serious		

© Assessment Instrument Document

Date generated: 24/10/2014 Page 40 of 40